

# Anesthetic Management in Patients with Renal Transplant Undergoing Coronary Bypass Graft Surgery

Gholam Ali Mollasadeghi, MD; Evaz Heidarpour, MD; Mohammad Hassan Kalantar Motamedi, MD\*; Rasool Farasatkish, MD; Kamal Raissi, MD

## Abstract

**Background-** By the end of 2000, more than 10,000 patients had received renal transplants in the Islamic Republic of Iran (IRI), and this number is expected to increase yearly. Since the 1-year survival rate for renal transplant recipients is approaching 90% and is continuing to improve annually, an increasing number of patients who have received renal transplants present for coronary artery bypass graft (CABG) surgery. They represent a technically demanding group of patients who require special consideration regarding preserving renal graft function and minimizing possible complications of cardiac surgery. This study was conducted to evaluate the outcome in renal transplant patients undergoing CABG surgery.

**Patients and Methods-** We prospectively studied seventeen renal transplant patients with approximately normal preoperative renal function (plasma creatinine 1.1-1.5 mg/dL) scheduled for elective coronary artery bypass surgery. Various aspects of anesthesia and the surgical procedures were assessed as regards the function of the transplanted kidney.

**Results-** Renal blood flow and renal transplant function are influenced before the induction of anesthesia, after sternotomy and before cardiopulmonary bypass (CPB), during hypothermic CPB and normothermic CPB, after sternal closure, and postoperative bleeding.

**Conclusion-** Protective interventions are very important in renal transplant patients undergoing coronary artery bypass graft surgery to prevent deterioration of renal function. Hemofiltration was performed routinely to prevent volume overload and excessive hemodilution. We used low-dose dopamine infusion (renal dose) throughout the operation and phenylephrine infusion during cardiopulmonary bypass. (*Iranian Heart Journal*. 2002; 2(4)&3(1): 14-20)

**Key words:** renal transplanted patients ■ coronary bypass graft ■ immunosuppressive drugs ■ kidney protection

With advances in anesthetic and surgical management, a larger number of sick patients, including many with renal dysfunction and renal transplants are being referred for cardiac surgery.<sup>1-4</sup> Particularly of importance is the

fact that coronary artery disease is the second most common cause of mortality in renal transplant patients, following renal factors such as graft rejection. Preoperative renal dysfunction and renal transplantation are both well-known predictors of

From the Department of Cardiac Anesthesia, Shaheed Rajaie Cardiovascular Medical Center, Mellat Park, Vali Asr Avenue, Tehran, Iran  
\*Dept. of Cardiovascular Surgery, Jamaran Heart Hospital, Baqiatallah University of Medical Sciences, Tehran, Iran  
Correspondence to: G. Mollasadeghi, MD, Shaheed Rajaie Cardiovascular Medical Center, Mellat Park, Vali Asr Avenue, Tehran, Iran  
Email: [roknabad@rhc.ac.ir](mailto:roknabad@rhc.ac.ir) Fax: + (9821) 2042026

postoperative renal failure, infection and renal rejection, which remain a serious postoperative complication of surgical procedures, especially those in which cardiopulmonary bypass (CPB) are utilized. Transplant recipients are always under treatment with various regimens of immunosuppressive drug therapy. The immunosuppressive drugs in common use are cyclosporine A, azathioprine, anti-lymphocyte globulin, monoclonal antibodies, and steroids. Newer drugs, such as tacrolimus (FK 506), may replace cyclosporin, and the drug mycophenolate (mofetil) may replace azathioprine in some clinical protocols.<sup>5</sup>

Because cyclosporine or tacrolimus plasma levels must be kept within the indicated therapeutic range, the blood levels of these agents in those renal transplanted patients receiving these drugs should be monitored daily in the perioperative period. Clinically significant reductions of cyclosporine or tacrolimus blood levels can be caused by dilution with massive fluid infusion preoperatively<sup>6</sup> and cardiopulmonary bypass.<sup>7</sup> Extracorporeal circulation may have a negative impact on renal function. Generally however, there is no consensus regarding the etiological factors for this detrimental effect.<sup>12</sup>

Our studies have shown that patients with normally functioning preoperative renal transplants do not suffer detrimental effects from extracorporeal circulation under moderately hypothermic bypass.<sup>3</sup> This study was carried out to evaluate the factors influencing immediate outcome in patients with renal transplants and with an acceptable renal function who undergo coronary artery bypass surgery and hemodilution and moderately hypothermic CPB.

### Methods and Materials

This study was performed prospectively on seventeen patients with normally functioning renal transplants, defined as

plasma creatinine levels between 1.1-1.5 mg/dL, who were scheduled for elective coronary artery bypass surgery. They ranged from 38 to 61 years of age (mean  $50.20 \pm 7.10$ ), had no recent myocardial infarctions, and had not been exposed to nephrotoxic drugs; including radiographic contrast media, for at least 3 days. Emergency coronary surgery, unstable anginal pain in the preoperative period, hemodynamic instability, and the use of inotropes or diuretics were considered as exclusion criteria.

### Anesthetic/operative technique

The patients all received a continuous infusion of 3mg/kg/min dopamine from the time they arrived in the operating room until they were extubated postoperatively. Invasive monitoring was completed before the induction of anesthesia and included left radial artery cannulation, central venous pressure, arterial blood gases analysis, oropharyngeal and rectal temperatures, pulse oximetry, and a Foley catheter for urinary output measurement.<sup>8</sup> Anesthetic management was uniform in all the patients. Premedication consisted of 0.1mg/kg morphine IM and 1mg/kg promethazine orally 30 min. preoperatively and 10mg diazepam orally the night before the operation. The induction of anesthesia was done with 20mg/kg sodium pentobarbital within one minute, midazolam 5mg, pancuronium bromide 0.12mg/kg and after 3 minutes of pre-oxygenation with pure oxygen via face mask, oro-tracheal intubation was performed with a suitable endotracheal tube. The maintenance of anesthesia was achieved with the infusion of midazolam, pancuronium bromide and fentanyl, without any volatile anesthetic agents or nitrous oxide. A nitroglycerine infusion of 1.5mg/kg/min was given to all the patients. After induction, 10-15mg/kg of Ringer's solution was infused to maintain filling pressures near the baseline and a cardiac index above 2.2 L/min/m<sup>2</sup>. The urinary

output was measured and if it was less than 1 ml/kg/h before CPB, an additional 5 ml/kg Ringer's solution was given. If the urinary output was below 2 ml/kg/h during CPB or less than 1 ml/kg/h after CPB, 10 mg furosemide was administered intravenously. Two patients had hypotension and tachycardia and received 3- $\mu$ g/kg/min epinephrine, and dopaminergic doses of dopamine, and their blood pressure and heart rate returned to the acceptable range.

Non-pulsatile extracorporeal circulation was performed with a roller pump and a microporous hollow fiber membrane oxygenator, primed with 1.5 L of Ringer's solution and 0.5-1g/kg of mannitol. The pump flow was at an average of 2.2 L/min/m<sup>2</sup>.

Moderate hypothermia (28-32° C) was used in all the patients. The hematocrit level was not allowed to descend below 25%. A hollow fiber hemofilter was utilized throughout the period the patients were on cardiopulmonary bypass.

Blood samples were obtained and analyzed at the following times: 1) pre-anesthetic measurement before the induction, 2) after the induction and at the beginning of mechanical ventilation, 3) during early hypothermic extracorporeal circulation, 4) during normothermic bypass, 5) during sternal closure, 6) 1 hour into the postoperative period. During the perioperative period, urine was collected in plastic graduated bottles.

The surgical procedure was carried out as routine for CABG surgery, paying particular attention to meticulous skin prep and total asepsis, and considering the immunosuppressed state of the patients due to chronic immunosuppressive therapy.

## Results

From the 17 patients who underwent coronary bypass surgery in this study, 16 patients recovered without any complication after surgery and were discharged from hospital within 2 weeks after the operation. One patient with severe left ventricular dysfunction (preoperative ejection fraction 20%) died a few hours after surgery because of intractable cardiac failure. The revascularization was technically difficult in these patients because of advanced coronary and peripheral atherosclerosis.

The renal function values of these patients during their stay in hospital were within the normal range, and no instances of increased blood urea and nitrogen levels or creatinine levels were seen (Table I).